

University of Groningen

**Efficient Havinga–Kondepudi resolution of conglomerate amino acid derivatives by slow cooling and abrasive grinding**

Leeman, Michel; Noorduyn, Wim L.; Millemaggi, Alessia; Vlieg, Elias; Meekes, Hugo; Enckevort, Willem J.P. van; Kaptein, Bernard; Kellogg, Richard M.

*Published in:*  
 CrystEngComm

*DOI:*  
 [10.1039/c0ce00140f](https://doi.org/10.1039/c0ce00140f)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
 Publisher's PDF, also known as Version of record

*Publication date:*  
 2010

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Leeman, M., Noorduyn, W. L., Millemaggi, A., Vlieg, E., Meekes, H., Enckevort, W. J. P. V., Kaptein, B., & Kellogg, R. M. (2010). Efficient Havinga–Kondepudi resolution of conglomerate amino acid derivatives by slow cooling and abrasive grinding. *CrystEngComm*, 12(7), 2051-2053. <https://doi.org/10.1039/c0ce00140f>

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

# Supplementary Material (ESI) for CrystEngComm  
# This journal is (c) The Royal Society of Chemistry 2010

```
data_global
_journal_name_full          CrystEngComm

_journal_codен_Cambridge    1350
_journal_volume             ?
_journal_page_first         ?
_journal_year               ?

_publ_contact_author_name   R.M.Kellogg
_publ_contact_author_email  r.m.kellogg@syncom.nl

_publ_section_title
;
Efficient Havinga-Kondepudi Resolution of Conglomerate
Amino Acid Derivatives by Slow Cooling and Abrasive Grinding
;
_publ_contact_author_address
;Kadijk 3
9747 AT Groningen
The Netherlands
;
loop_
_publ_author_name
R.M.Kellogg
'Michel Leeman'
'Wim Noorduin'
'Elias Vlieg'
'Hugo Meekes'
;
W.van Enckevort
;
'Alessia Millmaggi'
'Bernard Kaptein'

# Attachment '- Leeman.cif'

data_leeman
_database_code_depnum_ccdc_archive 'CCDC 773977'
loop_

_audit_creation_method
SHELXL-97
_chemical_name_systematic
;
(S)-2-((2-methylbenzylidene)amino)-2-phenylacetamide
;
_chemical_name_common
(S)-2-((2-methylbenzylidene)amino)-2-phenylacetamide
_chemical_formula_moiety      ?
_chemical_formula_sum         'C16 H16 N2 O'
_chemical_formula_weight      252.31

loop_
_atom_type_symbol
_atom_type_description
_atom_type_scatter_dispersion_real
_atom_type_scatter_dispersion_imag
_atom_type_scatter_source
C C 0.0033 0.0016 'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'
H H 0.0000 0.0000 'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'
N N 0.0061 0.0033 'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'
O O 0.0106 0.0060 'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'
```

```

_symmetry_cell_setting      ?
_symmetry_space_group_name_H-M  ?

loop_
_symmetry_equiv_pos_as_xyz
'x, y, z'
'x+1/2, -y+1/2, -z'
'-x, y+1/2, -z+1/2'
'-x+1/2, -y, z+1/2'

_cell_length_a              5.2025(2)
_cell_length_b              9.7343(8)
_cell_length_c              26.324(2)
_cell_angle_alpha           90.00
_cell_angle_beta            90.00
_cell_angle_gamma           90.00
_cell_volume                1333.10(16)
_cell_formula_units_Z       4
_cell_measurement_temperature 208(2)
_cell_measurement_reflns_used ?
_cell_measurement_theta_min  ?
_cell_measurement_theta_max  ?

_exptl_crystal_description  ?
_exptl_crystal_colour       ?
_exptl_crystal_size_max     ?
_exptl_crystal_size_mid     ?
_exptl_crystal_size_min     ?
_exptl_crystal_density_meas ?
_exptl_crystal_density_diffrn 1.257
_exptl_crystal_density_method 'not measured'
_exptl_crystal_F_000         536
_exptl_absorpt_coefficient_mu 0.080
_exptl_absorpt_correction_type ?
_exptl_absorpt_correction_T_min ?
_exptl_absorpt_correction_T_max ?

_exptl_special_details
;
?
;

_diffrn_ambient_temperature 208(2)
_diffrn_radiation_wavelength 0.71073
_diffrn_radiation_type       MoK\alpha
_diffrn_radiation_source     'fine-focus sealed tube'
_diffrn_radiation_monochromator graphite
_diffrn_measurement_device_type ?
_diffrn_measurement_method   ?
_diffrn_detector_area_resol_mean ?
_diffrn_standards_number     ?
_diffrn_standards_interval_count ?
_diffrn_standards_interval_time ?
_diffrn_standards_decay_%    ?
_diffrn_reflns_number         15101
_diffrn_reflns_av_R_equivalents 0.0321
_diffrn_reflns_av_sigmaI/netI 0.0436
_diffrn_reflns_limit_h_min    -6
_diffrn_reflns_limit_h_max    5
_diffrn_reflns_limit_k_min    -12
_diffrn_reflns_limit_k_max    12
_diffrn_reflns_limit_l_min    -34
_diffrn_reflns_limit_l_max    34
_diffrn_reflns_theta_min      2.23
_diffrn_reflns_theta_max      27.50

```

```

_reflns_number_total      2947
_reflns_number_gt         2232
_reflns_threshold_expression >2sigma(I)

_computing_data_collection ?
_computing_cell_refinement ?
_computing_data_reduction ?
_computing_structure_solution 'SHELXS-97 (Sheldrick, 1990)'
_computing_structure_refinement 'SHELXL-97 (Sheldrick, 1997)'
_computing_molecular_graphics ?
_computing_publication_material ?

_refine_special_details
;
Refinement of F2 against ALL reflections. The weighted R-factor wR and
goodness of fit S are based on F2, conventional R-factors R are based
on F, with F set to zero for negative F2. The threshold expression of
F2 > 2sigma(F2) is used only for calculating R-factors(gt) etc. and is
not relevant to the choice of reflections for refinement. R-factors based
on F2 are statistically about twice as large as those based on F, and R-
factors based on ALL data will be even larger.
;

_refine_ls_structure_factor_coef Fsqd
_refine_ls_matrix_type full
_refine_ls_weighting_scheme
'calc w=1/[\s2(Fo2)+(0.0341P)2+0.2820P] where P=(Fo2+2Fc2)/3'
_refine_ls_solution_primary direct
_refine_ls_solution_secondary difmap
_refine_ls_solution_hydrogens geom
_refine_ls_hydrogen_treatment mixed
_refine_ls_extinction_method none
_refine_ls_extinction_coef ?
_refine_ls_abs_structure_details 'Flack H D (1983), Acta Cryst. A39, 876-881'
_refine_ls_abs_structure_Flack 0.3(18)
_refine_ls_number_reflns 2947
_refine_ls_number_parameters 179
_refine_ls_number_restraints 0
_refine_ls_R_factor_all 0.0794
_refine_ls_R_factor_gt 0.0478
_refine_ls_wR_factor_ref 0.0922
_refine_ls_wR_factor_gt 0.0829
_refine_ls_goodness_of_fit_ref 1.088
_refine_ls_restrained_S_all 1.088
_refine_ls_shift/su_max 0.000
_refine_ls_shift/su_mean 0.000

loop_
_atom_site_label
_atom_site_type_symbol
_atom_site_fract_x
_atom_site_fract_y
_atom_site_fract_z
_atom_site_U_iso_or_equiv
_atom_site_adp_type
_atom_site_occupancy
_atom_site_calc_flag
_atom_site_refinement_flags
_atom_site_disorder_assembly
_atom_site_disorder_group
C1 C 0.4764(3) 0.45010(19) 0.11181(7) 0.0236(4) Uani 1 d . . .
H1 H 0.6357 0.5053 0.1092 0.028 Uiso 1 calc R . .
N2 N 0.2583(3) 0.54517(16) 0.11676(6) 0.0263(4) Uani 1 d . . .
C3 C 0.3013(4) 0.66571(19) 0.10142(6) 0.0277(4) Uani 1 d . . .
H3 H 0.4619 0.6841 0.0865 0.033 Uiso 1 calc R . .
C4 C 0.4543(4) 0.36310(18) 0.06414(6) 0.0230(4) Uani 1 d . . .

```

O5 O 0.6520(2) 0.31253(14) 0.04642(5) 0.0317(3) Uani 1 d . . .  
 N6 N 0.2252(3) 0.34532(17) 0.04435(6) 0.0298(4) Uani 1 d . . .  
 H6A H 0.093(4) 0.381(2) 0.0602(8) 0.036 Uiso 1 d . . .  
 H6B H 0.204(4) 0.296(2) 0.0165(8) 0.036 Uiso 1 d . . .  
 C11 C 0.4984(4) 0.35925(19) 0.15805(6) 0.0249(4) Uani 1 d . . .  
 C12 C 0.7023(4) 0.3759(2) 0.19084(8) 0.0386(5) Uani 1 d . . .  
 H12 H 0.8264 0.4439 0.1844 0.046 Uiso 1 calc R . .  
 C13 C 0.7250(5) 0.2931(3) 0.23314(8) 0.0539(7) Uani 1 d . . .  
 H13 H 0.8632 0.3062 0.2556 0.065 Uiso 1 calc R . .  
 C14 C 0.5494(5) 0.1926(3) 0.24267(8) 0.0515(7) Uani 1 d . . .  
 H14 H 0.5670 0.1362 0.2714 0.062 Uiso 1 calc R . .  
 C15 C 0.3469(5) 0.1740(2) 0.21011(9) 0.0507(6) Uani 1 d . . .  
 H15 H 0.2262 0.1041 0.2164 0.061 Uiso 1 calc R . .  
 C16 C 0.3201(4) 0.2577(2) 0.16814(8) 0.0381(5) Uani 1 d . . .  
 H16 H 0.1793 0.2456 0.1462 0.046 Uiso 1 calc R . .  
 C21 C 0.1174(4) 0.77857(19) 0.10532(7) 0.0267(5) Uani 1 d . . .  
 C22 C 0.1305(4) 0.8896(2) 0.07173(8) 0.0326(5) Uani 1 d . . .  
 C23 C -0.0334(5) 0.9993(2) 0.07945(9) 0.0431(6) Uani 1 d . . .  
 H23 H -0.0270 1.0747 0.0572 0.052 Uiso 1 calc R . .  
 C24 C -0.2062(5) 1.0004(2) 0.11914(9) 0.0450(6) Uani 1 d . . .  
 H24 H -0.3140 1.0768 0.1239 0.054 Uiso 1 calc R . .  
 C25 C -0.2215(4) 0.8906(2) 0.15168(8) 0.0405(5) Uani 1 d . . .  
 H25 H -0.3399 0.8910 0.1786 0.049 Uiso 1 calc R . .  
 C26 C -0.0609(4) 0.7799(2) 0.14430(7) 0.0316(5) Uani 1 d . . .  
 H26 H -0.0727 0.7038 0.1662 0.038 Uiso 1 calc R . .  
 C27 C 0.3141(5) 0.8913(2) 0.02752(9) 0.0478(6) Uani 1 d . . .  
 H27A H 0.4893 0.8951 0.0401 0.072 Uiso 1 calc R . .  
 H27B H 0.2802 0.9711 0.0065 0.072 Uiso 1 calc R . .  
 H27C H 0.2911 0.8085 0.0075 0.072 Uiso 1 calc R . .

loop\_

\_atom\_site\_aniso\_label

\_atom\_site\_aniso\_U\_11

\_atom\_site\_aniso\_U\_22

\_atom\_site\_aniso\_U\_33

\_atom\_site\_aniso\_U\_23

\_atom\_site\_aniso\_U\_13

\_atom\_site\_aniso\_U\_12

C1 0.0215(10) 0.0234(10) 0.0259(9) -0.0004(8) -0.0001(8) 0.0029(8)  
 N2 0.0285(9) 0.0225(8) 0.0278(8) -0.0020(7) -0.0001(7) 0.0032(7)  
 C3 0.0271(11) 0.0294(11) 0.0265(9) -0.0007(8) -0.0028(8) 0.0000(9)  
 C4 0.0260(11) 0.0211(10) 0.0220(8) 0.0003(8) 0.0015(8) 0.0019(8)  
 O5 0.0265(7) 0.0376(8) 0.0311(7) -0.0079(6) 0.0037(6) 0.0067(7)  
 N6 0.0237(9) 0.0378(10) 0.0278(8) -0.0102(8) -0.0025(7) 0.0023(8)  
 C11 0.0285(11) 0.0235(10) 0.0228(8) -0.0046(8) 0.0037(8) 0.0069(9)  
 C12 0.0400(13) 0.0420(13) 0.0337(10) -0.0020(9) -0.0071(10) 0.0020(11)  
 C13 0.0579(17) 0.0695(17) 0.0343(11) 0.0051(12) -0.0128(12) 0.0147(15)  
 C14 0.0658(17) 0.0590(16) 0.0296(11) 0.0157(11) 0.0056(12) 0.0232(15)  
 C15 0.0575(16) 0.0476(14) 0.0471(13) 0.0158(12) 0.0134(12) 0.0044(13)  
 C16 0.0386(13) 0.0412(12) 0.0344(11) 0.0032(10) 0.0023(9) 0.0004(11)  
 C21 0.0295(11) 0.0226(11) 0.0279(10) -0.0021(8) -0.0085(8) -0.0005(8)  
 C22 0.0318(12) 0.0273(11) 0.0386(11) 0.0040(9) -0.0082(9) -0.0013(9)  
 C23 0.0491(14) 0.0247(11) 0.0554(14) 0.0064(10) -0.0117(12) 0.0027(11)  
 C24 0.0453(14) 0.0285(12) 0.0612(14) -0.0085(11) -0.0054(13) 0.0134(11)  
 C25 0.0409(13) 0.0376(12) 0.0431(12) -0.0089(10) 0.0003(10) 0.0094(11)  
 C26 0.0382(12) 0.0240(11) 0.0327(10) -0.0006(9) -0.0040(9) -0.0011(9)  
 C27 0.0510(15) 0.0425(13) 0.0499(13) 0.0149(11) 0.0032(11) -0.0006(12)

\_geom\_special\_details

;

All esds (except the esd in the dihedral angle between two l.s. planes)  
 are estimated using the full covariance matrix. The cell esds are taken  
 into account individually in the estimation of esds in distances, angles  
 and torsion angles; correlations between esds in cell parameters are only  
 used when they are defined by crystal symmetry. An approximate (isotropic)  
 treatment of cell esds is used for estimating esds involving l.s. planes.

;

```
loop_  
_geom_bond_atom_site_label_1  
_geom_bond_atom_site_label_2  
_geom_bond_distance  
_geom_bond_site_symmetry_2  
_geom_bond_publ_flag  
C1 N2 1.470(2) . ?  
C1 C11 1.509(3) . ?  
C1 C4 1.518(2) . ?  
N2 C3 1.261(2) . ?  
C3 C21 1.461(3) . ?  
C4 O5 1.232(2) . ?  
C4 N6 1.312(2) . ?  
C11 C12 1.377(3) . ?  
C11 C16 1.381(3) . ?  
C12 C13 1.380(3) . ?  
C13 C14 1.362(3) . ?  
C14 C15 1.370(3) . ?  
C15 C16 1.380(3) . ?  
C21 C26 1.383(3) . ?  
C21 C22 1.398(3) . ?  
C22 C23 1.381(3) . ?  
C22 C27 1.506(3) . ?  
C23 C24 1.378(3) . ?  
C24 C25 1.372(3) . ?  
C25 C26 1.378(3) . ?
```

```
loop_  
_geom_angle_atom_site_label_1  
_geom_angle_atom_site_label_2  
_geom_angle_atom_site_label_3  
_geom_angle  
_geom_angle_site_symmetry_1  
_geom_angle_site_symmetry_3  
_geom_angle_publ_flag  
N2 C1 C11 110.86(14) . . ?  
N2 C1 C4 111.46(15) . . ?  
C11 C1 C4 110.22(14) . . ?  
C3 N2 C1 114.87(16) . . ?  
N2 C3 C21 124.14(18) . . ?  
O5 C4 N6 123.73(16) . . ?  
O5 C4 C1 118.19(16) . . ?  
N6 C4 C1 118.07(17) . . ?  
C12 C11 C16 118.76(18) . . ?  
C12 C11 C1 119.66(18) . . ?  
C16 C11 C1 121.58(17) . . ?  
C11 C12 C13 120.2(2) . . ?  
C14 C13 C12 120.7(2) . . ?  
C13 C14 C15 119.7(2) . . ?  
C14 C15 C16 120.0(2) . . ?  
C15 C16 C11 120.6(2) . . ?  
C26 C21 C22 119.65(18) . . ?  
C26 C21 C3 119.90(17) . . ?  
C22 C21 C3 120.34(17) . . ?  
C23 C22 C21 118.3(2) . . ?  
C23 C22 C27 119.81(19) . . ?  
C21 C22 C27 121.86(19) . . ?  
C24 C23 C22 121.3(2) . . ?  
C25 C24 C23 120.3(2) . . ?  
C24 C25 C26 119.1(2) . . ?  
C25 C26 C21 121.23(19) . . ?
```

```
_diffraction_measured_fraction_theta_max 0.988  
_diffraction_reflections_theta_full 25.00
```

_diffrn_measured_fraction_theta_full	0.996
_refine_diff_density_max	0.175
_refine_diff_density_min	-0.194
_refine_diff_density_rms	0.043